

POST-OPERATIVE INSTRUCTIONS FOR CYSTOSCOPIC PROCEDURES

Important- Please Read Completely

1. It is normal to have some burning or frequent urination after this procedure. Pyridium (Phenazopyridine) or ProceD EC taken as directed should help with this. It normally turns the urine orange or blue. If this hasn't been prescribed or you run out of the medicine, contact our office so that we can call your pharmacy.
2. If you have more than a little blood in your urine or on your undergarments, call the doctor. We advise no heavy lifting, exertion, or sexual activity for 1-2 weeks after this procedure. If constipation is experienced, a laxative of your choice and/or stool softener is highly recommended.
3. If antibiotics were prescribed, take as directed. Call the doctor for fever, chills, or any unusual pain in the abdomen, back, or genitals.
4. If a foley catheter was left in your bladder, it is normal to have some pain or discomfort in your abdomen or genitals. Take the pain pills (Lortab, Percocet, Darvocet N-100) as prescribed. If this doesn't control your discomfort, call the doctor.
5. Someone from our staff will call you within 24 hours of your surgery or after being discharged from the hospital to check on your condition. If you haven't been contacted within this time frame, please contact our office as soon as possible to schedule your follow-up appointment. You may be asked to come in to have your catheter removed with the nurses followed by a separate post-operative visit with the doctor.